

— ABLYBIO, Help Your Research



TCF7L2 Rabbit mAb

货号: **AYM29408**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50 FC: 1:20 - 1:50
理论分子量	68kDa
实测分子量	68kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SW620,HeLa,U-251MG,HepG2
细胞定位	nucleoplasm,nucleus,PML body
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human TCF7L2.
------	---


靶点信息

研究背景	This gene encodes a high mobility group (HMG) box-containing transcription factor that plays a key role in the Wnt signaling pathway. The protein has been implicated in blood glucose homeostasis. Genetic variants of this gene are associated with increased risk of type 2 diabetes. Several transcript variants encoding multiple different isoforms have been found for this gene.[provided by RefSeq, Oct 2010]
基因ID	6934
基因名	TCF7L2
Swiss	Q9NQB0 (https://www.uniprot.org/uniprotkb/Q9NQB0/entry)
别名	TCF7L2,TCF7L2 Rabbit mAb,HMG box transcription factor 4,T-cell-specific transcription factor 4,TCF4

产品验证



Western blot analysis of TCF7L2 expressed in SW620, HeLa, U-251MG using TCF7L2 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.



Immunohistochemical analysis of paraffin-embedded human colonic cancer, using TCF7L2 Antibody.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)