

Radixin Rabbit mAb

货号: **AYM29410**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50 FC: 1:20 - 1:50
理论分子量	80kDa
实测分子量	80kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,HepG2,Jurkat,C6,Mouse lung,Mouse kidney,Rat lung,Rat liver
细胞定位	adherens junction,apical plasma membrane,cortical actin cytoskeleton,extracellular exosome,extracellular space,microvillus,plasma membrane,T-tubule
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human Radixin.
------	--

靶点信息


研究背景	Radixin is a cytoskeletal protein that may be important in linking actin to the plasma membrane. It is highly similar in sequence to both ezrin and moesin. The radixin gene has been localized by fluorescence in situ hybridization to 11q23. A truncated version representing a pseudogene (RDXP2) was assigned to Xp21.3. Another pseudogene that seemed to lack introns (RDXP1) was mapped to 11p by Southern and PCR analyses. Multiple alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, May 2012]
------	---

基因ID	5962
基因名	RDX
Swiss	P35241
别名	Radixin

产品验证



Western blot analysis of Radixin expressed in Mouse lung, Rat lung, HeLa using Radixin Rabbit m Ab at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins : 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.



Immunohistochemical analysis of paraffin-embedded human spleen, using Radixin Antibody.

实验步骤

访问官网浏览详情: www.ablybio.cn