

Ezrin Rabbit mAb

货号: **AYM29443**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	80kDa
实测分子量	80kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	HCT116
细胞定位	Apical cell membrane, Cell projection, Cytoplasm, Cytoplasmic side, Peripheral membrane protein, cell cortex, cytoskeleton, microvillus membrane, ruffle membrane
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human Ezrin.
序列	IEVQQMKAQAREEKHQKQLERQQLTEKKRRETVEREKEQMMREKEELMLRLQDYEEKTKKAERELSEQIQRALQLEEK RAQEEAERLEADRMAALRAK

靶点信息

研究背景	The cytoplasmic peripheral membrane protein encoded by this gene functions as a protein-tyrosine kinase substrate in microvilli. As a member of the ERM protein family, this protein serves as an intermediate between the plasma membrane and the actin cytoskeleton. This protein plays a key role in cell surface structure adhesion, migration and organization, and it has been implicated in various human cancers. A pseudogene located on chromosome 3 has been identified for this gene. Alternatively spliced variants have also been described for this gene. [provided by RefSeq, Jul 2008]
------	---

基因ID	7430
基因名	EZR
Swiss	P15311
别名	CVIL;CVL;HEL-S-105;VIL2;EZR;Ezrin;ezrin

产品验证



Western blot analysis of Ezrin expressed in HCT116 using Ezrin Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn