

# RAB7 Rabbit mAb

货号: **AYM29463**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC FC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200 <b>IF/ICC:</b> 1:50 - 1:200 <b>FC:</b> 1:20 - 1:50
理论分子量	23kDa
实测分子量	23kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,Neuro-2a,C2C12,Mouse brain,PC-12,Rat brain
细胞定位	Cytoplasmic side,Cytoplasmic vesicle,Late endosome membrane,Lipid droplet,Lysosome membrane,Melanosome membrane,Peripheral membrane protein,autophagosome membrane,phagosome membrane
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein corresponding to Human RAB7.
------	---------------------------------------------------------

## 靶点信息


研究背景	RAB family members are small, RAS-related GTP-binding proteins that are important regulators of vesicular transport. Each RAB protein targets multiple proteins that act in exocytic / endocytic pathways. This gene encodes a RAB family member that regulates vesicle traffic in the late endosomes and also from late endosomes to lysosomes. This encoded protein is also involved in the cellular vacuolation of the VacA cytoxin of Helicobacter pylori. Mutations at highly conserved amino acid residues in this gene have caused some forms of Charcot-Marie-Tooth (CMT) type 2 neuropathies.
基因ID	7879

基因名	RAB7A
Swiss	P51149
别名	RAB7

## 产品验证



Western blot analysis of RAB7 expressed in Mouse brain, Rat brain, HeLa using RAB7 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30 ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.



Immunohistochemical analysis of paraffin-embedded human colon, using RAB7 Antibody.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)