

MUC2 Rabbit mAb

货号: **AYM29468**

产品信息

反应	Human,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50 FC: 1:20 - 1:50
理论分子量	540kDa
实测分子量	540kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HT-29,Mouse small intestine,Rat large intestine,Rat small intestine
细胞定位	Secreted
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human MUC2.
序列	SELQTEGRTRNHGHNV CSTWGNFHYKTFDGDVFRFPGLCDYNFASDCRGSYKEFAVHLKRGPGQAEAPAGVESILLTIK DDTIYLRHLAVLNGAVVSTPHYSPLLLIEKSDAYTKVYSRAGLTLMWNREDALMLELDTKFRNHTCGLCGDYNGLSYSE FLSDGVLFSPLEFGNMQKIN

靶点信息

研究背景	This gene encodes a member of the mucin protein family. Mucins are high molecular weight glycoproteins produced by many epithelial tissues. The protein encoded by this gene is secreted and forms an insoluble mucous barrier that protects the gut lumen. The protein polymerizes into a gel of which 80% is composed of oligosaccharide side chains by weight. The protein features a central domain containing tandem repeats rich in threonine and proline that varies between 50 and 115 copies in different individuals. Alternatively spliced transcript variants of this gene have been described, but their full-length nature is not known.
基因ID	4583
基因名	
Swiss	Q02817
别名	MUC2;MLP;MUC-2;SMUC;mucin-2

产品验证



Western blot analysis of MUC2 expressed in Rat large intestine, HT-29, Mouse small intestine using MUC2 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn