

BAT3 Rabbit mAb

货号: **AYM29476**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	96kDa/113kDa/118kDa/119kDa/122kDa
实测分子量	160kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,HeLa,Mouse testis,Rat testis
细胞定位	Cytoplasm,Nucleus,cytosol
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human BAT3.
序列	MEPNDSTSTAVEEPDSLEVLVKTLDTSQTRTFIVGAQMNVKEFKEHIAASVSIPSEKQRLIYQGRVLQDDKKLQEYNVGGKVI HLVERAPPQTHLPSGASSGTGSASATHGGGSPPGTRGPGASVHDRNANSYVMVGTFLPSDGSADVHINMEQAPIQS EPRVRLVMAQHMIRDIQTLLSRMECRGGPQPQHSQPPQPQ

靶点信息

研究背景	This gene was first characterized as part of a cluster of genes located within the human major histocomp atibility complex class III region. This gene encodes a nuclear protein that is cleaved by caspase 3 and is i mplicated in the control of apoptosis. In addition, the protein forms a complex with E1A binding protein p 300 and is required for the acetylation of p53 in response to DNA damage. Multiple transcript variants en coding different isoforms have been found for this gene.
------	--

基因ID	7917
基因名	BAG6
Swiss	P46379
别名	BAG6;BAG-6;BAT3;D6S52E;G3

产品验证



Western blot analysis of BAT3 expressed in Mouse testis,Rat testis,293T using BAT3 Rabbit mAb at 1:10 00. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Bl ocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn