

BCKDK Rabbit mAb

货号: **AYM29478**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	37kDa/41kDa/46kDa
实测分子量	46kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,HepG2,PC-12,Mouse pancreas
细胞定位	Mitochondrion,Mitochondrion matrix
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human BCKDK.
------	--

靶点信息

研究背景	The branched-chain alpha-ketoacid dehydrogenase complex (BCKD) is an important regulator of the valine, leucine, and isoleucine catabolic pathways. The protein encoded by this gene is found in the mitochondrion, where it phosphorylates and inactivates BCKD. Several transcript variants encoding different isoforms have been found for this gene.
基因ID	10295
基因名	BCKDK
Swiss	O14874

别名	BCKDK
----	-------

产品验证



Western blot analysis of BCKDK expressed in Mouse pancreas, HeLa, HepG2 using BCKDK Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn