

— ABLYBIO, Help Your Research



FADD Rabbit mAb

货号: **AYM29519**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IP FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IP: 1:20 - 1:50 FC: 1:20 - 1:50
理论分子量	23kDa
实测分子量	28kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,MCF7,HeLa,Mouse liver,Mouse kidney,Mouse heart
细胞定位	cytoplasm,cytosol,neuron projection,nucleus,plasma membrane,riposome
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human FADD .
------	--

靶点信息

研究背景	The protein encoded by this gene is an adaptor molecule that interacts with various cell surface receptors and mediates cell apoptotic signals. Through its C-terminal death domain, this protein can be recruited by TNFRSF6/Fas-receptor, tumor necrosis factor receptor, TNFRSF25, and TNFSF10/TRAIL-receptor, and thus it participates in the death signaling initiated by these receptors. Interaction of this protein with the receptors unmask the N-terminal effector domain of this protein, which allows it to recruit caspase-8, and thereby activate the cysteine protease cascade. Knockout studies in mice also suggest the importance of this protein in early T cell development.
基因ID	8772
基因名	FADD
Swiss	Q13158 (https://www.uniprot.org/uniprotkb/Q13158/entry)
别名	FADD,FADD Rabbit mAb,FAS-associating death domain-containing protein,Growth-inhibiting gene 3 protein,Mediator of receptor induced toxicity,MORT1

产品验证



Western blot analysis of FADD expressed in 293T,MCF7,HeLa using FADD Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)