

# CYP26A1 Rabbit mAb

货号: **AYM29521**

## 产品信息

反应	Human,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200
理论分子量	48kDa/56kDa
实测分子量	50kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2
细胞定位	
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein corresponding to Human CYP26A1.
序列	GFPFFGETLQMVLQRRKFLQMKRRKYGFYKTHLFGRPTVRVMGADNVRRILLGEHRLVSVHWPASVRTILGSGCLSNLH DSSHKQRKKVIMRAFSREALE

## 靶点信息

研究背景	This gene encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This endoplasmic reticulum protein acts on retinoids, including all-trans-retinoic acid (RA), with both 4-hydroxylation and 18-hydroxylation activities. This enzyme regulates the cellular level of retinoic acid which is involved in regulation of gene expression in both embryonic and adult tissues. Two alternatively spliced transcript variants of this gene, which encode the distinct isoforms, have been reported.
------	---

基因ID	1592
基因名	CYP26A1
Swiss	O43174
别名	CYP26A1;CP26;CYP26;P450RAI;P450RAI1

## 产品验证



Western blot analysis of CYP26A1 expressed in HepG2 using CYP26A1 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)