

# CD7 Rabbit mAb

货号: **AYM29545**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC ICC IP
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200 <b>ICC:</b> 1:50 - 1:200 <b>IP:</b> 1:20 - 1:50
理论分子量	25kDa
实测分子量	37kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat,U-937,Rat spleen
细胞定位	Membrane,Single-pass type I membrane protein
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein corresponding to Human CD7.
序列	AQEVQQSPHCTTVPVGASVNITCSTSGGLRGIYLRQLGPPQDIIYYEDGVVPTDRRFRGRIDFSGSQDNLITMHLQLSDTGTYTCAITEVNVYSGTLVLTVEEQSQGWHRCSDAPPRASALPAPPTGSALPDPQTASALPDPPAASALP

## 靶点信息

研究背景	This gene encodes a transmembrane protein which is a member of the immunoglobulin superfamily. This protein is found on thymocytes and mature T cells. It plays an essential role in T-cell interactions and also in T-cell/B-cell interaction during early lymphoid development.
基因ID	924

基因名	CD7
Swiss	P09564
别名	CD7;GP40;LEU-9;TP41;Tp40

## 产品验证



Western blot analysis of CD7 expressed in Jurkat,U-937,Rat spleen using CD7 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)