

RAB27A Rabbit mAb

货号: **AYM29561**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC IP FC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50 FC: 1:20 - 1:50
理论分子量	24kDa
实测分子量	28kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG(KD)
细胞定位	Late endosome,Lipid-anchor,Lysosome,Melanosome,Membrane
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human RAB27A.
序列	MSDGDYDYLIKFLALGDSGVGKTSVLYQYTDGKFNKFITTVGIDFREKRVVYRASGPDGATGRGQRIHLQLWDTAGQE RFRSLTTAFFRDAMGFLLLFDLTNEQSFLNVRNWISQLQMHAECENPDIVLCGNKSDLEDQRVVKEEEAIALAEKYGIPYFE TSAANGTNISQAIEMLLDLIMKRMERCVDKSWIPEGVVRSNGHASTDQLSEEKEKGACGC

靶点信息

研究背景	The protein encoded by this gene belongs to the small GTPase superfamily, Rab family. The protein is membrane-bound and may be involved in protein transport and small GTPase mediated signal transduction. Mutations in this gene are associated with Griscelli syndrome type 2. Alternative splicing occurs at this locus and four transcript variants encoding the same protein have been identified.
------	--

基因ID	5873
基因名	RAB27A
Swiss	P51159
别名	RAB27A;GS2;HsT18676;RAB27;RAM

产品验证



Western blot analysis of RAB27A expressed in U-87MG(KD) using RAB27A Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn