

— ABLYBIO, Help Your Research



c-Jun Rabbit mAb

货号: **AYM29569**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC ICC IP
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 ICC: 1:50 - 1:200 IP: 1:20 - 1:50
理论分子量	35kDa
实测分子量	43kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,HT-1080,NIH/3T3
细胞定位	Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human c-Jun.
------	--


靶点信息

研究背景	This gene is the putative transforming gene of avian sarcoma virus 17. It encodes a protein which is highly similar to the viral protein, and which interacts directly with specific target DNA sequences to regulate gene expression. This gene is intronless and is mapped to 1p32-p31, a chromosomal region involved in both translocations and deletions in human malignancies.
基因ID	3725
基因名	JUN
Swiss	P05412 (https://www.uniprot.org/uniprotkb/P05412/entry)
别名	c-Jun,c-Jun Rabbit mAb,JUN,Activator protein 1,Proto-oncogene c-Jun,Transcription factor AP-1 subunit Jun,V-jun avian sarcoma virus 17 oncogene homolog,p39


产品验证




Western blot analysis of c-Jun expressed in HeLa, HT-1080, NIH/3T3 using c-Jun Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.




Immunohistochemical analysis of paraffin-embedded Rat cerebral cortex, using the Antibody at 1:500 dilution.




Immunohistochemical analysis of paraffin-embedded Rat cerebellum, using the Antibody at 1:500 dilution.



Immunohistochemical analysis of paraffin-embedded Human squamous carcinoma, using the Antibody at 1:200 dilution.



Immunohistochemical analysis of paraffin-embedded Human breast cancer, using the Antibody at 1:200 dilution.



Immunohistochemical analysis of paraffin-embedded Mouse skeletal muscle - gastrocnemius, using the Antibody at 1:500 dilution.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)