

NSD3 Rabbit mAb

货号: **AYM29588**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB FC
推荐浓度	WB: 1:500 - 1:2000 FC: 1:20 - 1:50
理论分子量	72kDa/154kDa/156kDa/160kDa/161kDa
实测分子量	162kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T
细胞定位	Chromosome,Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human NSD3.
序列	EEALSQAKKSVASKTEVKKTRRPRSVLNTQPEQTNAGEVASSLSSTEIRRHSQRRHTSAEEEEPPPVKIAWKTAARKSLPA SITMHKGSLLDLQKCNMSPVVKIEQVFALQATGDGKFIDQFVYSTKGIGNKTEISVRGQDRLLIISTPNQRNEKPTQSVSSPE ATSGSTGSVEKKQRRSIRTRSESEKSTEVPKIK

靶点信息

研究背景	This gene is related to the Wolf-Hirschhorn syndrome candidate-1 gene and encodes a protein with PWWP (proline-tryptophan-tryptophan-proline) domains. This protein methylates histone H3 at lysine residues 4 and 27, which represses gene transcription. Two alternatively spliced variants have been described.
基因ID	54904

基因名	NSD3
Swiss	Q9BZ95
别名	NSD3;KMT3F;KMT3G;WHISTLE;WHSC1L1;pp14328

产品验证



Western blot analysis of NSD3 expressed in 293T using NSD3 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn