

— ABLYBIO, Help Your Research



# MINA53 Rabbit mAb

货号: **AYM29595**

## 产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IF/ICC:</b> 1:50 - 1:200
理论分子量	53kDa
实测分子量	53kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	BxPC-3,Mouse testis,Rat testis
细胞定位	cytosol,nucleolus,nucleoplasm
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein corresponding to Human MINA53.
------	---

## 靶点信息

研究背景	MINA is a c-Myc (MYC; MIM 190080) target gene that may play a role in cell proliferation or regulation of cell growth. (Tsuneoka et al., 2002 [PubMed 12091391]; Zhang et al., 2005 [PubMed 15897898]).[supplied by OMIM, May 2008]
基因ID	84864
基因名	RIOX2
Swiss	Q8IU8 (https://www.uniprot.org/uniprotkb/Q8IU8/entry)
别名	MINA53,MINA53 Rabbit mAb,RIOX2,60S ribosomal protein L27a histidine hydroxylase,Bifunctional lysine-specific demethylase and histidyl-hydroxylase MINA,Histone lysine demethylase MINA,MYC-induced nuclear antigen,Mineral dust-induced gene protein

## 产品验证



Western blot analysis of MINA53 expressed in BxPC-3, Mouse testis, Rat testis using MINA53 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (https://www.ablybio.cn/www.ablybio.cn)