

# RhoGAP Rabbit mAb

货号: **AYM29604**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC FC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200 <b>FC:</b> 1:20 - 1:50
理论分子量	26kDa/27kDa/172kDa
实测分子量	190kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,Jurkat,293T,HepG2,PC-12,Mouse brain,Rat brain
细胞定位	Cytoplasm,Membrane,Peripheral membrane protein
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein corresponding to Human RhoGAP.
序列	EGLYRVSGNKTDQDNIQKQFDQDHNINLVSM EVT VNAVAGALKAFFADLPDPLIPYSLHPELLEAAKIPDKTERLHALKEIV KKFHPVNYDVFYVITHLNRVSQQHKINLMTADNLSICFWPTLMRPDFENREFLSTTKIHQSVVETFIQQCQFFFYNGEIVET TNIVAPPPPSNPGQLVEPMVPLQLPPPLQPQLIQPQLQTDPLGII

## 靶点信息

研究背景	Rho GTPase activating protein 5 negatively regulates RHO GTPases, a family which may mediate cytoskeleton changes by stimulating the hydrolysis of bound GTP. Two transcript variants encoding different isoforms have been found for this gene.
基因ID	394

基因名	ARHGAP5
Swiss	Q13017
别名	ARHGAP5;GFI2;RhoGAP5;p190-B;p190BRhoGAP

## 产品验证



Western blot analysis of RhoGAP expressed in Mouse brain,Rat brain,HeLa using RhoGAP Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)