

GC / VDBP Rabbit mAb

货号: **AYM29609**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	39kDa/52kDa/55kDa
实测分子量	53kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Human serum,Mouse serum,THP-1
细胞定位	Secreted
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human GC / VDBP.
序列	AMDVFVCTYFMPAAQLPELPDVELPTNKDVCDPGNTKVMKDYTFELSRRTHLPEVFLSKVLEPTLKS LGECDDVEDSTTCF NAKGPLLKKESSFIDKQGQELCADYSENTFTEYKKKLAERLKAKLPDATPTELAKLVNKHSDFASNCCSINSPPLYCDSEIDA ELKNIL

靶点信息

研究背景	The protein encoded by this gene belongs to the albumin gene family. It is a multifunctional protein found in plasma, ascitic fluid, cerebrospinal fluid and on the surface of many cell types. It binds to vitamin D and its plasma metabolites and transports them to target tissues. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.
------	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

基因ID	2638
基因名	GC
Swiss	P02774
别名	GC;DBP;DBP/GC;GRD3;Gc-MAF;GcMAF;HEL-S-51;VDBG;VDBP

产品验证



Western blot analysis of GC / VDBP expressed in Human serum, Mouse serum, THP-1 using GC / VDBP R abbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins : 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn