

— ABLYBIO, Help Your Research



# TCEA1 Rabbit mAb

货号: **AYM29620**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200 <b>IF/ICC:</b> 1:50 - 1:200 <b>IP:</b> 1:20 - 1:50
理论分子量	31kDa/33kDa
实测分子量	34kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,293T
细胞定位	Nucleus
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein corresponding to Human TCEA1.
------	--

## 靶点信息

研究背景	Necessary for efficient RNA polymerase II transcription elongation past template-encoded arresting sites. The arresting sites in DNA have the property of trapping a certain fraction of elongating RNA polymerases that pass through, resulting in locked ternary complexes. Cleavage of the nascent transcript by S-II allows the resumption of elongation from the new 3'-terminus.
基因ID	6917
基因名	TCEA1
Swiss	P23193 ( <a href="https://www.uniprot.org/uniprotkb/P23193/entry">https://www.uniprot.org/uniprotkb/P23193/entry</a> )
别名	TCEA1,TCEA1 Rabbit mAb,Transcription elongation factor S-II protein 1,Transcription elongation factor TFII S.o,GTF2S,TFIIS

## 产品验证



Western blot analysis of TCEA1 expressed in HeLa,293T using TCEA1 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/>[www.ablybio.cn](http://www.ablybio.cn))