

PAK2 Rabbit mAb

货号: **AYM29636**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	58kDa
实测分子量	61kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,SKOV3,Jurkat
细胞定位	Cytoplasm,Cytoplasm,Lipid-anchor,Membrane,Nucleus,perinuclear region
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human PAK2.
序列	PQAVLDVLKIFYDSNTVKQKYLSTFPPEKDGFPSTPALNAKGTEAPAVVTEEDDDEETAPPVIAPRPDHTKSIYTRSVIDPVPAPVGDSHVDGAAKSLDKQKKTMTDEEIMEKLRTIV

靶点信息

研究背景	The p21 activated kinases (PAK) are critical effectors that link Rho GTPases to cytoskeleton reorganization and nuclear signaling. The PAK proteins are a family of serine/threonine kinases that serve as targets for the small GTP binding proteins, CDC42 and RAC1, and have been implicated in a wide range of biological activities. The protein encoded by this gene is activated by proteolytic cleavage during caspase-mediated apoptosis, and may play a role in regulating the apoptotic events in the dying cell.
------	--

基因ID	5062
基因名	PAK2
Swiss	Q13177
别名	PAK2;PAK65;PAKgamma

产品验证



Western blot analysis of PAK2 expressed in HeLa,SKOV3,Jurkat using PAK2 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn