

# RAP80 Rabbit mAb

货号: **AYM29637**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200 <b>IF/ICC:</b> 1:50 - 1:200
理论分子量	17kDa/38kDa/61kDa/70kDa/79kDa
实测分子量	80kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	MCF7,293T,Mouse thymus,Rat brain
细胞定位	Nucleus
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein corresponding to Human RAP80 .
序列	MPRRKKKVKEVSESRNLEKKDVETSSVSVKRRRLEDAFIVISDSGDGEEPKEENGLQKTKTKQSNRAKCLAKRKIAQMTE EEQFALALKMSEQEAREVNSQEEEEELLRKAIAESLNSCRPSDASATRSRPLATGPSSQSHQEKTDS

## 靶点信息

研究背景	This gene encodes a nuclear protein that interacts with Brca1 (breast cancer 1) in a complex to recognize and repair DNA lesions. This protein binds ubiquitinated lysine 63 of histone H2A and H2AX. This protein may also function as a repressor of transcription. Alternative splicing results in multiple transcript variant S.
基因ID	51720

基因名	UIMC1
Swiss	Q96RL1
别名	UIMC1;RAP80;X2HRIP110

## 产品验证



Western blot analysis of RAP80 expressed in MCF7,293T,Mouse thymus using RAP80 Rabbit mAb at 1:1 000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. B locking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)