

Nck2 Rabbit mAb

货号: **AYM29639**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 ICC: 1:50 - 1:200
理论分子量	42kDa
实测分子量	47kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,Mouse lung,Rat brain
细胞定位	Cytoplasm,Endoplasmic reticulum
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Nck2.
序列	MTEEVIVIAKWDYTAQQDQELDIKNERLWLLDDSKTWWRVNRNAANRTGYVPSNYVERKNSLKKGSLVKNLKDTLGLGK TRRKTSARDASPTPSTDAEYPANGSGADRIYDLNIPAFVKFAYVAEREDELVLKGSRTVMEKCSGDGWWRGSYNGQIG WFPSNYVLEEVDAAAESPFLSLRKGASLSNGQGSRVLHVVTLYPFSSVTEELNFEKGETMEVIEKPENDPEWWKCK NARGQVGLVPKNYVVVLSLSDGPALHPAHAPQISYTGPSSSGRFAGREWYYGNVTRHQAECALNERGVEGDFLIRDSESSP SDFSVSLKASGKNKHFVKQLVDNVYCIGQRRFHTMDELVEHYKKAPIFTSEHGKLYLVRALQ

靶点信息

研究背景	This gene encodes a member of the NCK family of adaptor proteins. The protein contains three SH3 domains and one SH2 domain. The protein has no known catalytic function but has been shown to bind and recruit various proteins involved in the regulation of receptor protein tyrosine kinases. It is through these regulatory activities that this protein is believed to be involved in cytoskeletal reorganization. Alternate transcriptional splice variants, encoding different isoforms, have been characterized.
基因ID	8440
基因名	NCK2
Swiss	O43639
别名	NCK2;GRB4;NCKbeta

产品验证



Western blot analysis of Nck2 expressed in 293T, Mouse lung, Rat brain using Nck2 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn