

— ABLYBIO, Help Your Research



# GTF2F2 Rabbit mAb

货号: **AYM29673**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200 <b>IF/ICC:</b> 1:50 - 1:200 <b>IP:</b> 1:20 - 1:50
理论分子量	28kDa
实测分子量	30kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2,Mouse thymus,Mouse skeletal muscle
细胞定位	Nucleus
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein corresponding to Human GTF2F2.
------	---

## 靶点信息

研究背景	TFIIIF is a general transcription initiation factor that binds to RNA polymerase II and helps to recruit it to the initiation complex in collaboration with TFIIB. It promotes transcription elongation. This subunit shows ATP-dependent DNA-helicase activity.
基因ID	2963
基因名	GTF2F2
Swiss	P13984 ( <a href="https://www.uniprot.org/uniprotkb/P13984/entry">https://www.uniprot.org/uniprotkb/P13984/entry</a> )
别名	GTF2F2,GTF2F2 Rabbit mAb,General transcription factor IIF 30 kDa subunit,Transcription initiation factor IIF subunit beta,Transcription initiation factor RAP30,RAP30

## 产品验证



Western blot analysis of GTF2F2 expressed in HepG2, Mouse thymus, Mouse skeletal muscle using GTF2F2 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/>[www.ablybio.cn](http://www.ablybio.cn))