

— ABLYBIO, Help Your Research



# ADAMTS13 Rabbit mAb

货号: **AYM29684**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB FC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>FC:</b> 1:20 - 1:50
理论分子量	39kDa/144kDa/147kDa/153kDa
实测分子量	154kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2,LO2,22Rv1,U-251MG,Mouse liver
细胞定位	Secreted
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein corresponding to Human ADAMTS13.
------	---

## 靶点信息

研究背景	This gene encodes a member of a family of proteins containing several distinct regions, including a metalloproteinase domain, a disintegrin-like domain, and a thrombospondin type 1 (TS) motif. The enzyme encoded by this gene specifically cleaves von Willebrand Factor (vWF). Defects in this gene are associated with thrombotic thrombocytopenic purpura. Alternative splicing results in multiple transcript variants.
基因ID	11093
基因名	ADAMTS13
Swiss	Q76LX8 ( <a href="https://www.uniprot.org/uniprotkb/Q76LX8/entry">https://www.uniprot.org/uniprotkb/Q76LX8/entry</a> )
别名	ADAMTS13,ADAMTS13 Rabbit mAb,von Willebrand factor-cleaving protease,C9orf8

## 产品验证



Western blot analysis of ADAMTS13 expressed in HepG2, LO2, 22Rv1 using ADAMTS13 Rabbit mAb at 1:10 00. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/>[www.ablybio.cn](http://www.ablybio.cn))