

# ADAMTS13 Rabbit mAb

货号: **AYM29684**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB FC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>FC:</b> 1:20 - 1:50
理论分子量	39kDa/144kDa/147kDa/153kDa
实测分子量	154kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2,LO2,22Rv1,U-251MG,Mouse liver
细胞定位	Secreted
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein corresponding to Human ADAMTS13.
序列	LSPGAPLKRPPSPGFQRQRQRRAAGGILHLELLVAVGPDVFAQHQEDTERYVLTNLNIGAELLRDPGLGAQFRVHLVK MVILTEPEGAPNITANLTSSLLSVCWWSQTINPEDDTPGHADLVLYITRFDLELPDGNRQVRGVTQLGGACSPWWSCLITE DTGFDLGVITIAHEIGHSFGLHEDGAPSGCGPSGHVMSDGAAPRAGLAWSPCSRRQLLSLLSAGRARCWVDPWRPQP GSAGHPPDAQ

## 靶点信息

研究背景	This gene encodes a member of a family of proteins containing several distinct regions, including a metalloproteinase domain, a disintegrin-like domain, and a thrombospondin type 1 (TS) motif. The enzyme encoded by this gene specifically cleaves von Willebrand Factor (vWF). Defects in this gene are associated with thrombotic thrombocytopenic purpura. Alternative splicing results in multiple transcript variants.
------	--

基因ID	11093
基因名	ADAMTS13
Swiss	Q76LX8
别名	ADAMTS13;ADAM-TS13;ADAMTS-13;C9orf8;VWF-CP

## 产品验证



Western blot analysis of ADAMTS13 expressed in HepG2, LO2, 22Rv1 using ADAMTS13 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)