

RAB5C / RABL Rabbit mAb

货号: **AYM29701**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 ICC: 1:50 - 1:200
理论分子量	23kDa/27kDa
实测分子量	25kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-549,BT-474,HL-60,Mouse lung
细胞定位	Cell membrane,Cytoplasmic side,Early endosome membrane,Lipid-anchor,Melanosome
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human RAB5C / RABL.
序列	MAGRGGGAARPNNGPAAGNKICQFKLVLLGESAVGKSSLVLRVFKGQFHEYQESTIGAAFLTQTVCCLDDTTVKFEIWDTAGQ ERYHSLAPMYRGAQAIVVYDITNTDTFARAKNWVKELQRQASPNIVIALAGNKADLASKRAVEFQEAQAYADDNSLLFM ETSAKTAMNVNEIFMAIAKKLPKNEPQNATGAPGRNRGVLDLQENNPASRSQCCSN

靶点信息

研究背景	Members of the Rab protein family are small GTPases of the Ras superfamily that are thought to ensure fidelity in the process of docking and/or fusion of vesicles with their correct acceptor compartment (Han et al., 1996 [PubMed 8646882]).
基因ID	5878

基因名	RAB5C
Swiss	P51148
别名	RAB5C;L1880;RAB5CL;RAB5L;RABL

产品验证



Western blot analysis of RAB5C /L expressed in Mouse lung,A-549,BT-474 using RAB5C / RABL Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30u g per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn