

GABARAP Rabbit mAb

货号: **AYM29716**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	17kDa
实测分子量	17kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,293T+Chloroquine,NIH/3T3,NIH/3T3+Chloroquine,C6,C6+Chloroquine
细胞定位	Cytoplasm,Cytoplasmic vesicle,Endomembrane system,Golgi apparatus membrane,autophagosome,cyto skeleton
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human GABARAP.
------	--

靶点信息

研究背景	Gamma-aminobutyric acid A receptors [GABA(A) receptors] are ligand-gated chloride channels that mediate inhibitory neurotransmission. This gene encodes GABA(A) receptor-associated protein, which is highly positively charged in its N-terminus and shares sequence similarity with light chain-3 of microtubule-associated proteins 1A and 1B. This protein clusters neurotransmitter receptors by mediating interaction with the cytoskeleton. [provided by RefSeq, Jul 2008]
基因ID	11337

基因名	GABARAP
Swiss	O95166
别名	GABARAP

产品验证



Western blot analysis of GABARAP expressed in 293T,293T+Chloroquine,NIH/3T3 using GABARAP Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 1 20s.

实验步骤

访问官网浏览详情: www.ablybio.cn