

U1A Rabbit mAb

货号: **AYM29737**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC IP
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50
理论分子量	31kDa
实测分子量	31kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat,293T,A-431,MCF7,Mouse brain,Mouse testis,Mouse thymus,Rat brain
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human U1A.
序列	MAVPETRPNHTIYINNLLNEKIKKDELKKSLEYAIFSQFGQILDILVSRSLKMRGQAFVIFKEVSSATNALRSMQGFPPYDKPMRI QYAKTDSDIIAKMGTFVERDRKREKRKPKSQETPATKKAVQGGGATPVVGAVQGPVPGMPPMTQAPRIMHHMPGQPP YMPPPGMIPPPGLAPGQIPPGAMPPQQLMPGQMPPAQPLSENPPNHILFTNLPEETNELMLSMLFNQFPGFKEVRLVPGR HDIAFVEFDNEVQAGAARDALQGFKITQNNAMKISFAKK

靶点信息

研究背景	The protein encoded by this gene associates with stem loop II of the U1 small nuclear ribonucleoprotein, which binds the 5' splice site of precursor mRNAs and is required for splicing. The encoded protein autoregulates itself by polyadenylation inhibition of its own pre-mRNA via dimerization and has been implicated in the coupling of splicing and polyadenylation.
------	---

基因ID	6626
基因名	SNRPA
Swiss	P09012
别名	SNRPA;Mud1;U1-A;U1A

产品验证



Western blot analysis of U1A expressed in Mouse brain,Rat brain,Jurkat using U1A Rabbit mAb at 1:1000 . Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn