

# Vitronectin Rabbit mAb

货号: **AYM29777**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200 <b>IF/ICC:</b> 1:50 - 1:200
理论分子量	54kDa
实测分子量	75kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Human serum,Rat serum
细胞定位	Secreted,extracellular space
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein corresponding to Human Vitronectin .
序列	SLAKKQRFRRNRKGYRSQRGHSRGRNQNSRRPSRATWLSLFSSEESNLGANNYDDYRMDWLVPATCEPIQSVFFFSG DKYYRVNLRTRRVDTVDPYPYRSIAQYWLGCAPAGHL

## 靶点信息

研究背景	The protein encoded by this gene is a member of the pexin family. It is found in serum and tissues and promotes cell adhesion and spreading, inhibits the membrane-damaging effect of the terminal cytolytic complement pathway, and binds to several serpin serine protease inhibitors. It is a secreted protein and exists in either a single chain form or a clipped, two chain form held together by a disulfide bond.
基因ID	7448

基因名	VTN
Swiss	P04004
别名	VTN;V75;VN;VNT

## 产品验证



Western blot analysis of Vitronectin expressed in Human serum,Rat serum using Vitronectin Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)