

— ABLYBIO, Help Your Research



# Zic1 Rabbit mAb

货号: **AYM29784**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC FC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IF/ICC:</b> 1:50 - 1:200 <b>FC:</b> 1:20 - 1:50
理论分子量	
实测分子量	48kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG,U-251MG,Mouse brain,Rat brain
细胞定位	cytoplasm,nucleoplasm,nucleus
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Zic1.
------	---

## 靶点信息

研究背景	This gene encodes a member of the ZIC family of C2H2-type zinc finger proteins. Members of this family are important during development. Aberrant expression of this gene is seen in medulloblastoma, a childhood brain tumor. This gene is closely linked to the gene encoding zinc finger protein of the cerebellum 4, a related family member on chromosome 3. This gene encodes a transcription factor that can bind and transactivate the apolipoprotein E gene. [provided by RefSeq, Jul 2008]
基因ID	7545
基因名	ZIC1
Swiss	Q15915 ( <a href="https://www.uniprot.org/uniprotkb/Q15915/entry">https://www.uniprot.org/uniprotkb/Q15915/entry</a> )
别名	Zic1,Zic1 Rabbit mAb,Zinc finger protein 201,Zinc finger protein of the cerebellum 1,ZIC,ZNF201

## 产品验证



Western blot analysis of Zic1 expressed in Mouse brain,Rat brain,U-87MG using Zic1 Rabbit mAb at 1:100. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/www.ablybio.cn>)