

— ABLYBIO, Help Your Research



SRP19 Rabbit mAb

货号: **AYM29815**

产品信息

| | |
|-------|---|
| 反应 | Human,Mouse,Rat |
| 宿主 | Rabbit |
| 克隆性 | Monoclonal |
| 预测反应 | |
| 应用 | WB IHC IF/ICC |
| 推荐浓度 | WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 |
| 理论分子量 | 8kDa/12kDa/16kDa |
| 实测分子量 | 20kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | HepG2,MCF7 |
| 细胞定位 | Cytoplasm |
| 纯化 | Affinity purification |


抗原信息

| | |
|------|--|
| 抗原信息 | Recombinant fusion protein corresponding to Human SRP19. |
|------|--|

靶点信息

| | |
|-------|--|
| 研究背景 | Signal-recognition-particle assembly, binds directly to 7S RNA and mediates binding of the 54 kDa subunit of the SRP. |
| 基因ID | 6728 |
| 基因名 | SRP19 |
| Swiss | P09132 (https://www.uniprot.org/uniprotkb/P09132/entry) |
| 别名 | SRP19,SRP19 Rabbit mAb |

产品验证



Western blot analysis of SRP19 expressed in HepG2, MCF7 using SRP19 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)