

DCR2 Rabbit mAb

货号: **AYM29819**

产品信息

| | |
|-------|-----------------------------------------------------------------------------------------------|
| 反应 | Human |
| 宿主 | Rabbit |
| 克隆性 | Monoclonal |
| 预测反应 | |
| 应用 | WB IHC IF/ICC |
| 推荐浓度 | WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 |
| 理论分子量 | 41kDa |
| 实测分子量 | 36kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | THP-1 |
| 细胞定位 | Membrane,Single-pass type I membrane protein |
| 纯化 | Affinity purification |

抗原信息


| | |
|------|---------------------------------------------------------------------------------------------------------------------------------------|
| 抗原信息 | Recombinant fusion protein corresponding to Human DCR2. |
| 序列 | VLFRRRSCPSRVPGAEDNARNETLSNRYLQPTQVSEQEIQGQELAEELTGVTVLPEEPQRLLEQAEAEAGCQRRRLVNVND ADSADISTLLDASATLEEGHAKETIQDQLVGSEKLFYEDEAGSATSCL |

靶点信息

| | |
|------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 研究背景 | The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor contains an extracellular TRAIL-binding domain, a transmembrane domain, and a truncated cytoplamic death domain . This receptor does not induce apoptosis, and has been shown to play an inhibitory role in TRAIL-induced cell apoptosis. |
| 基因ID | 8793 |

| | |
|-------|----------------------------------------------|
| 基因名 | TNFRSF10D |
| Swiss | Q9UBN6 |
| 别名 | TNFRSF10D;CD264;DCR2;TRAIL-R4;TRAILR4;TRUNDD |

产品验证



Western blot analysis of DCR2 expressed in THP-1 using DCR2 Rabbit mAb at 1:1000. Secondary antibody : HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn