

— ABLYBIO, Help Your Research



# DCR2 Rabbit mAb

货号: **AYM29819**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200 <b>IF/ICC:</b> 1:50 - 1:200
理论分子量	41kDa
实测分子量	36kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	THP-1
细胞定位	Membrane,Single-pass type I membrane protein
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein corresponding to Human DCR2.
------	---

## 靶点信息

研究背景	The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor contains an extracellular TRAIL-binding domain, a transmembrane domain, and a truncated cytoplasmic death domain. This receptor does not induce apoptosis, and has been shown to play an inhibitory role in TRAIL-induced cell apoptosis.
基因ID	8793
基因名	TNFRSF10D
Swiss	Q9UBN6 ( <a href="https://www.uniprot.org/uniprotkb/Q9UBN6/entry">https://www.uniprot.org/uniprotkb/Q9UBN6/entry</a> )
别名	DCR2,DCR2 Rabbit mAb,TNFRSF10D,Decoy receptor 2,TNF-related apoptosis-inducing ligand receptor 4,TRAIL receptor with a truncated death domain,TRAILR4,TRUNDD

## 产品验证



Western blot analysis of DCR2 expressed in THP-1 using DCR2 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/www.ablybio.cn>)