

HDGF Rabbit mAb

货号: **AYM29820**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	25kDa/26kDa/28kDa
实测分子量	37kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	22Rv1,HeLa,A-549,293T,U-251MG,MCF7,BT-474
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human HDGF.
序列	MSRSNRQKEYKCGDLVFAKMKGYPHWPARIDEMPEAAVKSTANKYQVFFFGTHETAFLGPKDLFPYEEESKEKFGKPNKR KGFSEGLWEIENNPVKASGYQSSQKSCVEEPEPEPEAAEGDGDKGNAEGSSDEEGKLVIDEPAKEKNEKGALKRRA GDILLEDSPKRPKEAENPEGEEKEAATLEVERPLMEVEKKNSTPSEPGSGRPPQEEEEEEDEEEEATKEDAEAPGIRDHESL

靶点信息

研究背景	This gene encodes a member of the hepatoma-derived growth factor family. The encoded protein has mitogenic and DNA-binding activity and may play a role in cellular proliferation and differentiation. High level s of expression of this gene enhance the growth of many tumors. This gene was thought initially to be located on chromosome X; however, that location has been determined to correspond to a related pseudogene. Alternatively spliced transcript variants encoding distinct isoforms have been described.
------	--

基因ID	3068
基因名	HDGF
Swiss	P51858
别名	HDGF;HMG1L2

产品验证



Western blot analysis of HDGF expressed in 22Rv1, HeLa, A-549 using HDGF Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn