

— ABLYBIO, Help Your Research



Torsin A Rabbit mAb

货号: **AYM29823**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB FC
推荐浓度	WB: 1:500 - 1:2000 FC: 1:20 - 1:50
理论分子量	22kDa/37kDa
实测分子量	35kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	MCF7,BT-474,DU145
细胞定位	Cell projection,Cytoplasm,Cytoplasmic vesicle,Cytoplasmic vesicle membrane,Endoplasmic reticulum lumen,Nucleus membrane,Peripheral membrane protein,cytoskeleton,growth cone,secretory vesicle,synaptic vesicle
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Torsin A.
------	-------------------------------------------------------------

靶点信息

研究背景	The protein encoded by this gene is a member of the AAA family of adenosine triphosphatases (ATPases), is related to the Clp protease/heat shock family and is expressed prominently in the substantia nigra pars compacta. Mutations in this gene result in the autosomal dominant disorder, torsion dystonia 1.
基因ID	1861
基因名	TOR1A
Swiss	O14656 (https://www.uniprot.org/uniprotkb/O14656/entry)
别名	Torsin A,Torsin A Rabbit mAb,TOR1A,Dystonia 1 protein,Torsin ATPase-1A,Torsin family 1 member A,DQ2,DYT1,TA,TORA

产品验证



Western blot analysis of Torsin A expressed in MCF7,BT-474,DU145 using Torsin A Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)