— ABLYBIO, Help Your Research

Myb Rabbit mAb

货号: AYM29832

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IP ICC/IF
推荐浓度	**WB:** 1:500 - 1:2000 **IHC:** 1:50 - 1:200 **IF/ICC:** 1:50 - 1:200 **IP:** 1:20 - 1:50
理论分子量	37-46kDa/63-85kDa
实测分子量	80kDa,72kDa
形式	Liquid
保存条件	Store at -20℃. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	无明确组织表达
细胞定位	Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Myb.


靶点信息

研究背景	This gene encodes a protein with three HTH DNA-binding domains that functions as a transcription regula tor. This protein plays an essential role in the regulation of hematopoiesis. This gene may be aberrently e xpressed or rearranged or undergo translocation in leukemias and lymphomas, and is considered to be a n oncogene. Alternative splicing results in multiple transcript variants.
基因ID	4602
基因名	MYB
Swiss	P10242 (https://www.uniprot.org/uniprotkb/P10242/entry)
别名	Myb,Myb Rabbit mAb,Proto-oncogene c-Myb

产品验证



Western blot analysis of Myb expressed in human esophagus cancer using Myb Rabbit mAb at 1:1000. Se condary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking bu ffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.



Immunofluorescence analysis of MCF-7 cells labelling Myb with . The cells were fixed with cold 100% met hanol (10min, 4℃) and blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween 20 fo r 1h. The cells were then incubate with Goat Anti-Rabbit IgG Fluor488 (1:200，shown in green) at room te mprature for 1h.

实验步骤

访问官网浏览详情: www.ablybio.cn (https://www.ablybio.cn//www.ablybio.cn)