

— ABLYBIO, Help Your Research



UBE2C Rabbit mAb

货号: **AYM29852**

产品信息

| | |
|-------|---|
| 反应 | Human |
| 宿主 | Rabbit |
| 克隆性 | Monoclonal |
| 预测反应 | |
| 应用 | WB |
| 推荐浓度 | WB: 1:500 - 1:2000 |
| 理论分子量 | 15kDa/16kDa/17kDa/19kDa |
| 实测分子量 | 20kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | PC-3,NIH/3T3 |
| 细胞定位 | cytosol,nucleoplasm,nucleus,plasma membrane |
| 纯化 | Affinity purification |


抗原信息

| | |
|------|--|
| 抗原信息 | Recombinant fusion protein corresponding to Human UBE2C. |
|------|--|

靶点信息

| | |
|-------|--|
| 研究背景 | The modification of proteins with ubiquitin is an important cellular mechanism for targeting abnormal or short-lived proteins for degradation. Ubiquitination involves at least three classes of enzymes: ubiquitin-activating enzymes, ubiquitin-conjugating enzymes, and ubiquitin-protein ligases. This gene encodes a member of the E2 ubiquitin-conjugating enzyme family. The encoded protein is required for the destruction of mitotic cyclins and for cell cycle progression, and may be involved in cancer progression. Multiple transcript variants encoding different isoforms have been found for this gene. Pseudogenes of this gene have been defined on chromosomes 4, 14, 15, 18, and 19. |
| 基因ID | 11065 |
| 基因名 | UBE2C |
| Swiss | O00762 (https://www.uniprot.org/uniprotkb/O00762/entry) |
| 别名 | UBE2C,UBE2C Rabbit mAb,(E3-independent) E2 ubiquitin-conjugating enzyme C,E2 ubiquitin-conjugating enzyme C,UbcH10,Ubiquitin carrier protein C,Ubiquitin-protein ligase C |

产品验证



Western blot analysis of UBE2C expressed in PC-3,NIH/3T3 using UBE2C Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)