

GRK2 Rabbit mAb

货号: **AYM29858**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	79kDa
实测分子量	80kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HT-29,Mouse spleen,Mouse brain
细胞定位	Cell membrane,Cytoplasm
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human GRK2.
序列	LGRGAQEVKESPFRRSLDWQMVFLQKYPPLIPRGEVNAADAFDIGSFDEEDTKGIKLLDSDQELYRNFPLTISERWQQE VAETVFDTINAETDRLEARKKAKNKQLGHEEDYALGKDCIMHGYMSKMGNPFLTQWQRRYFYLFNRLWRGEGEAPQS LLTMEEIQSVEETQIKERKCLLLKIRGGKQFILQCSDPELVQWKELRDAYREAQQLVQRVPMKNKPRSPVVELSKVPLV QRGSANGL

靶点信息

研究背景	The product of this gene phosphorylates the beta-2-adrenergic receptor and appears to mediate agonist-specific desensitization observed at high agonist concentrations. This protein is an ubiquitous cytosolic enzyme that specifically phosphorylates the activated form of the beta-adrenergic and related G-protein-coupled receptors. Abnormal coupling of beta-adrenergic receptor to G protein is involved in the pathogenesis of the failing heart.
------	---

基因ID	156
基因名	GRK2
Swiss	P25098
别名	GRK2;ADRBK1;BARK1;BETA-ARK1

产品验证



Western blot analysis of GRK2 expressed in HT-29, Mouse spleen, Mouse brain using GRK2 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane . Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn