

PARP Rabbit mAb

货号: **AYM29873**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	113kDa
实测分子量	113kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,Jurkat,A-549,Mouse testis,Rat testis
细胞定位	Nucleus,nucleolus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human PARP.
------	---

靶点信息


研究背景	This gene encodes a chromatin-associated enzyme, poly(ADP-ribosyl)transferase, which modifies various nuclear proteins by poly(ADP-ribosyl)ation. The modification is dependent on DNA and is involved in the regulation of various important cellular processes such as differentiation, proliferation, and tumor transfor mation and also in the regulation of the molecular events involved in the recovery of cell from DNA dama ge. In addition, this enzyme may be the site of mutation in Fanconi anemia, and may participate in the pa thophysiology of type I diabetes.
基因ID	142

基因名	PARP1
Swiss	P09874
别名	PARP

产品验证



Western blot analysis of PARP expressed in Mouse testis,Rat testis,HeLa using PARP Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.



Immunohistochemical analysis of paraffin-embedded human testis, using PARP Antibody.

实验步骤

访问官网浏览详情: www.ablybio.cn