

# S6K1 Rabbit mAb

货号: **AYM29889**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IP
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200 <b>IP:</b> 1:20 - 1:50
理论分子量	51kDa/52kDa/56kDa/59kDa
实测分子量	70kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	NIH/3T3
细胞定位	Cell junction,Cytoplasm,Mitochondrion outer membrane,Mitochondrion,Nucleus,synapse,synaptosome
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein corresponding to Human S6K1.
序列	GFTYV

## 靶点信息

研究背景	This gene encodes a member of the ribosomal S6 kinase family of serine/threonine kinases. The encoded protein responds to mTOR (mammalian target of rapamycin) signaling to promote protein synthesis, cell growth, and cell proliferation. Activity of this gene has been associated with human cancer. Alternatively spliced transcript variants have been observed. The use of alternative translation start sites results in isoforms with longer or shorter N-termini which may differ in their subcellular localizations. There are two pseudogenes for this gene on chromosome 17.
------	---

基因ID	6198
基因名	RPS6KB1
Swiss	P23443
别名	PS6K;S6K;S6K-beta-1;S6K1;STK14A;p70 S6KA;p70(S6K)-alpha;p70-S6K;p70-alpha;P70 S6K;RPS6KB1;p70S6KA

## 产品验证



Western blot analysis of S6K1 expressed in NIH/3T3 using S6K1 Rabbit mAb at 1:1000. Secondary antibody y: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)