

PRPF31 Rabbit mAb

货号: **AYM29891**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC FC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	30kDa/40kDa/54kDa/55kDa
实测分子量	60kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-251MG,Jurkat,HepG2,Mouse lung,Mouse testis,Rat lung,Rat testis
细胞定位	Cajal body,Nucleus,Nucleus speckle
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human PRPF31.
序列	MSLADELLADLEEAEEEEEGGSYGEEEEPAIEDVQEETQLDLSGDSVKTIAKLWDSKMFAEIMMKIEEYISKQAKASEVM GPVEAAPEYRVIVDANNLTVEIENELNIIHKFIRDKYSKRFPESLVPNALDYIRTVKELGNLSDKCKNNENLQILT NATIMV VSVTASTTQGQQLSEEEELERLEEACDMALELNAS

靶点信息

研究背景	This gene encodes a component of the spliceosome complex and is one of several retinitis pigmentosa-ca using genes. When the gene product is added to the spliceosome complex, activation occurs.
基因ID	26121

基因名	PRPF31
Swiss	Q8WWY3
别名	PRPF31;NY-BR-99;PRP31;RP11;SNRNP61

产品验证



Western blot analysis of PRPF31 expressed in U-251MG, Jurkat, HepG2 using PRPF31 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn