

RbAp48 Rabbit mAb

货号: **AYM29900**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50 FC: 1:20 - 1:50
理论分子量	43kDa/46kDa/47kDa
实测分子量	48kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T
细胞定位	Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human RbAp48.
序列	MADKEAAFDDAVEERVINEEYKIWKKNTPFLYDLVMTHALEWPSLTAQWLDPDVTRPEGKDFSIHRLVLGTHTSDEQNHLVI ASVQLPNDDAQFDASHYDSEKGEFGGFSVSGKIEIEIKINHEGEVNRARYMPQNPCIATKTPSSDVLVFDYTKHPSKPD PSGECNPDRLRLRGHQKEGYGLSWNPNLGHLSSASDDHTICLWDISAVPKEGKVVDKIFTGHTAVVEDVSWHLLHES LFGSVADDQKLMIWDRSNTSKPSHSVDAHTAEVNCLSFNPYSEFILATGSADKTVALWDLRNLKLLHSFESHKDEIF QVQWSPHNETILASSGTDRRLNVWDLKIGEEQSPEDAEDGPELFIHGGHTAKISDFSWNPNEPWWICSVSEDNIMQV WQMAENIYNDEDPEGSVDPEGQGS

靶点信息

研究背景	This gene encodes a ubiquitously expressed nuclear protein which belongs to a highly conserved subfamily of WD-repeat proteins. It is present in protein complexes involved in histone acetylation and chromatin assembly. It is part of the Mi-2 complex which has been implicated in chromatin remodeling and transcriptional repression associated with histone deacetylation. This encoded protein is also part of co-repressor complexes, which is an integral component of transcriptional silencing. It is found among several cellular proteins that bind directly to retinoblastoma protein to regulate cell proliferation. This protein also seems to be involved in transcriptional repression of E2F-responsive genes. Three transcript variants encoding different isoforms have been found for this gene.
基因ID	5928
基因名	RBBP4
Swiss	Q09028
别名	RBBP4;NURF55;RBAP48;lin-53

产品验证



Western blot analysis of RbAp48 expressed in 293T using RbAp48 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn