

— ABLYBIO, Help Your Research



NUP153 Rabbit mAb

货号: **AYM29903**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC FC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	149kDa/153kDa/157kDa
实测分子量	154kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse liver,Rat liver
细胞定位	Nucleus,Nucleus membrane,nuclear pore complex
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human NUP153.
------	---

靶点信息

研究背景	Nuclear pore complexes regulate the transport of macromolecules between the nucleus and cytoplasm. They are composed of at least 100 different polypeptide subunits, many of which belong to the nucleoporin family. Nucleoporins are glycoproteins found in nuclear pores and contain characteristic pentapeptide XFXFG repeats as well as O-linked N-acetylglucosamine residues oriented towards the cytoplasm. The protein encoded by this gene has three distinct domains: a N-terminal region containing a pore targeting and an RNA-binding domain, a central region containing multiple zinc finger motifs, and a C-terminal region containing multiple XFXFG repeats. Alternative splicing results in multiple transcript variants of this gene.
基因ID	9972
基因名	NUP153
Swiss	P49790 (https://www.uniprot.org/uniprotkb/P49790/entry)
别名	NUP153,NUP153 Rabbit mAb,153 kDa nucleoporin,Nucleoporin Nup153

产品验证



Western blot analysis of NUP153 expressed in Mouse liver,Rat liver using NUP153 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)