

U2AF65 Rabbit mAb

货号: **AYM29910**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	53kDa
实测分子量	54kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat,SW620,Mouse spleen
细胞定位	Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human U2AF65.
序列	NQDKNFLEFRSVDETTQAMAFDGIIFQGQSLKIRRPDYQPLPGMSENPSVYVPGVVSTVVPDSAHKLFIGGLPNYLN DDQVKELLTSFGPLKAFNLVKDSATGLSKGYAFCEYVDINVTDQAIAGLNGMQLGDKLLVQRASVGAKNATLSTINQTP VTLQVPGLMSSQVQMGGHPTEVLCLMNMVLPEELLDDDEEYEEIVEDVRDECSKYGLVKSIEIPRPVDGVEVPGCGKIFVEF TSVFDCQKAMQGLTGRKFANRVVVTKYCDPDSYHRRDFW

靶点信息

研究背景	U2 auxiliary factor (U2AF), comprised of a large and a small subunit, is a non-snRNP protein required for the binding of U2 snRNP to the pre-mRNA branch site. This gene encodes the U2AF large subunit which contains a sequence-specific RNA-binding region with 3 RNA recognition motifs and an Arg/Ser-rich domain necessary for splicing. The large subunit binds to the polypyrimidine tract of introns early during spliceosome assembly. Multiple transcript variants have been detected for this gene, but the full-length nature of only two have been determined to date.
基因ID	11338
基因名	U2AF2
Swiss	P26368
别名	U2AF2;U2AF65

产品验证



Western blot analysis of U2AF65 expressed in Jurkat,SW620,Mouse spleen using U2AF65 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn