

CITED2 Rabbit mAb

货号: **AYM29988**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	23kDa/28kDa
实测分子量	28kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SH-SY5Y,Mouse brain,Rat brain
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human CITED2.
序列	MSGLEMADHMMAMNHGRFPDGTNGLHHHPAHRMGMGQFSPHHHQQQPQHAFNALMGEHIIHYGAGNMNATSGI RHAMGPGTVNGGHPPSALAPAARFNNSQFMGPPVASQGGSLPASMQLQKLNNQYFNHHYPHNHYMPDLHPAAGHQ MNGTNQHFRDCNPKHSGGSSTPGGSGGSSTPGGSGSSGGGAGSSNSGGGSGSGNMPASVAHVPAAMLPPNVIDTD FIDEEVLMSLVIEMGLDRIKELPELWLGQNEFDFTDFVCKQQPSRVSC

靶点信息

研究背景	The protein encoded by this gene inhibits transactivation of HIF1A-induced genes by competing with binding of hypoxia-inducible factor 1-alpha to p300-CH1. Mutations in this gene are a cause of cardiac septal defects. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.
基因ID	10370

基因名	CITED2
Swiss	Q99967
别名	CITED2;ASD8;MRG-1;MRG1;P35SRJ;VSD2

产品验证



Western blot analysis of CITED2 expressed in SH-SY5Y, Mouse brain, Rat brain using CITED2 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn