

c-Fos Rabbit mAb

货号: **AYM29991**

产品信息

| | |
|-------|---|
| 反应 | Human,Mouse,Rat |
| 宿主 | Rabbit |
| 克隆性 | Monoclonal |
| 预测反应 | |
| 应用 | WB FC |
| 推荐浓度 | WB: 1:500 - 1:2000 FC: 1:20 - 1:50 |
| 理论分子量 | 28kDa/36kDa/40kDa |
| 实测分子量 | 62kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | HeLa |
| 细胞定位 | Cytoplasm,Endoplasmic reticulum,Nucleus,cytosol |
| 纯化 | Affinity purification |

抗原信息


| | |
|------|---|
| 抗原信息 | Recombinant fusion protein corresponding to Human c-Fos. |
| 序列 | MMFSGFNADYEASSSRCSASPAGDSLSEYHSPADSFSSMGSPVNAQDFCTDLAVSSANFIPTVTAISTSPDLQWLQPALVSSVAPSQTRAPHFPGVPAPSAGAYS RAGVVKT |

靶点信息

| | |
|------|---|
| 研究背景 | The Fos gene family consists of 4 members: FOS, FOSB, FOSL1, and FOSL2. These genes encode leucine zipper proteins that can dimerize with proteins of the JUN family, thereby forming the transcription factor complex AP-1. As such, the FOS proteins have been implicated as regulators of cell proliferation, differentiation, and transformation. In some cases, expression of the FOS gene has also been associated with apoptotic cell death. |
| 基因ID | 2353 |

| | |
|-------|--------------------------|
| 基因名 | FOS |
| Swiss | P01100 |
| 别名 | FOS;AP-1;C-FOS;p55;c-Fos |

产品验证



Western blot analysis of c-Fos expressed in HeLa using c-Fos Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn