— ABLYBIO, Help Your Research

c-Fos Rabbit mAb

货号: AYM29991

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB FC
推荐浓度	**WB:** 1:500 - 1:2000 **FC:** 1:20 - 1:50
理论分子量	28kDa/36kDa/40kDa
实测分子量	62kDa
形式	Liquid
保存条件	Store at -20℃. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa
细胞定位	Cytoplasm,Endoplasmic reticulum,Nucleus,cytosol
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human c-Fos.

靶点信息

研究背景	The Fos gene family consists of 4 members: FOS, FOSB, FOSL1, and FOSL2. These genes encode leucine zipper proteins that can dimerize with proteins of the JUN family, thereby forming the transcription factor complex AP-1. As such, the FOS proteins have been implicated as regulators of cell proliferation, different iation, and transformation. In some cases, expression of the FOS gene has also been associated with apo ptotic cell death.
基因ID	2353
基因名	FOS
Swiss	P01100 (https://www.uniprot.org/uniprotkb/P01100/entry)
别名	c-Fos,c-Fos Rabbit mAb,FOS,Cellular oncogene fos,Fos proto-oncogene,AP-1 transcription factor subunit,G 0/G1 switch regulatory protein 7,Proto-oncogene c-Fos,Transcription factor AP-1 subunit c-Fos,G0S7

产品验证



Western blot analysis of c-Fos expressed in HeLa using c-Fos Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dr y milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (https://www.ablybio.cn//www.ablybio.cn)