

Golgin 97 Rabbit mAb

货号: **AYM29995**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IP
推荐浓度	WB: 1:500 - 1:2000 IP: 1:20 - 1:50
理论分子量	88kDa
实测分子量	97kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	HepG2, HeLa, SW480, Mouse brain, Mouse heart, Mouse kidney, Mouse liver, Rat pancreas
细胞定位	Golgi apparatus, Golgi stack membrane, Peripheral membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human Golgin 97.
序列	MFAKLKKKIAEETAVAQRPGGATRIPRSVSKESVASMADSGDDFASDGSSSREDLSSQLLRNEQIRKLEARLSDYAEQVRNLQKIKEKLEIALEKHQDSSMRKFQEQNETFQANRAKMAEGLALALARKDQEWSEKMDQLEKEKNILTAQLQEMKNQSMNLFQRRDEMLEGFFQQELSKIKHMLLKKEESLGKMEQELEARRELSRTQEELMNSNQMSDDLQKLEELQRHYS TLEEQRDHVIASKTGAESKITALAQKEQELQALIQLSIDLQKVTAETQEKEDVITHLQEKV

靶点信息

研究背景	The Golgi apparatus, which participates in glycosylation and transport of proteins and lipids in the secretory pathway, consists of a series of stacked cisternae (flattened membrane sacs). Interactions between the Golgi and microtubules are thought to be important for the reorganization of the Golgi after it fragments during mitosis. This gene encodes one of the golgins, a family of proteins localized to the Golgi. This encoded protein is associated with Sjogren's syndrome.
------	--

基因ID	2800
基因名	GOLGA1
Swiss	Q92805
别名	GOLGA1;golgin-97;golgin A1

产品验证



Western blot analysis of Golgin 97 expressed in HepG2, HeLa, SW480 using Golgin 97 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn