— ABLYBIO, Help Your Research

Phospho-Histone H1.4 (T17) Rabbit mAb

货号: AYM30028

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC ICC/IF
推荐浓度	**WB:** 1:500 - 1:2000 **IHC:** 1:50 - 1:200 **IF/ICC:** 1:50 - 1:200
理论分子量	
实测分子量	30kDa
形式	Liquid
保存条件	Store at -20℃. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	无明确组织表达
细胞定位	
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human Phospho-Histone H1.4 (T17) .

靶点信息


研究背景	

基因ID	
基因名	
Swiss	P10412P16402 (https://www.uniprot.org/uniprotkb/P10412P16402/entry)
别名	Phospho-Histone H1.4 (T17),Phospho-Histone H1.4 (T17) Rabbit mAb


产品验证



Western blot analysis of Phospho-Histone H1.4 (T17) expressed in human kidney cancer,mouse heart,rat testis using Phospho-Histone H1.4 (T17) Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.



Immunohistochemical analysis of paraffin-embedded mouse colon, using Phospho-Histone H1.4 (T17) Antibody.



Immunofluorescence analysis of HeLa cells labelling Phospho-Histone H1.4 (T17) with . The cells were fixed with cold 100% methanol (10min, 4℃) and blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween 20 for 1h. The cells were then incubate with Goat Anti-Rabbit IgG Fluor488 (1:200，shown in green) at room temprature for 1h.

实验步骤

访问官网浏览详情: www.ablybio.cn (https://www.ablybio.cn//www.ablybio.cn)