

SNX1 Rabbit mAb

货号: **AYM30036**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	51kDa/59kDa/63kDa
实测分子量	74kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,Jurkat,LO2,A-549,HT-1080
细胞定位	Cell projection,Cytoplasmic side,Early endosome membrane,Endosome membrane,Golgi apparatus,Perip heral membrane protein,lamellipodium,trans-Golgi network membrane
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human SNX1.
------	---

靶点信息

研究背景	This gene encodes a member of the sorting nexin family. Members of this family contain a phox (PX) domain, which is a phosphoinositide binding domain, and are involved in intracellular trafficking. This endosomal protein regulates the cell-surface expression of epidermal growth factor receptor. This protein also has a role in sorting protease-activated receptor-1 from early endosomes to lysosomes. This protein may form oligomeric complexes with family members. This gene results in three transcript variants encoding distinct isoforms.
基因ID	6642

基因名	SNX1
Swiss	Q13596
别名	SNX1

产品验证



Western blot analysis of SNX1 expressed in HeLa, Jurkat, LO2 using SNX1 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn