— ABLYBIO, Help Your Research

Dnmt3a Rabbit mAb

货号: AYM30050

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC FC
推荐浓度	**WB:** 1:500 - 1:2000 **IHC:** 1:50 - 1:200 **IF/ICC:** 1:50 - 1:200 **FC:** 1:20 - 1:50
理论分子量	16kDa/81kDa/101kDa
实测分子量	130kDa
形式	Liquid
保存条件	Store at -20℃. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-549,293T,HeLa,HL-60,MCF7,HepG2,Mouse testis
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Dnmt3a.

靶点信息

研究背景	CpG methylation is an epigenetic modification that is important for embryonic development, imprinting, and X-chromosome inactivation. Studies in mice have demonstrated that DNA methylation is required for mammalian development. This gene encodes a DNA methyltransferase that is thought to function in de novo methylation, rather than maintenance methylation. The protein localizes to the cytoplasm and nucleus and its expression is developmentally regulated.
基因ID	1788
基因名	DNMT3A
Swiss	Q9Y6K1 (https://www.uniprot.org/uniprotkb/Q9Y6K1/entry)
别名	Dnmt3a,Dnmt3a Rabbit mAb,Cysteine methyltransferase DNMT3A,DNA methyltransferase HsaIIIA

产品验证



Western blot analysis of Dnmt3a expressed in Mouse testis,A-549,293T using Dnmt3a Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (https://www.ablybio.cn//www.ablybio.cn)