

USP9x Rabbit mAb

货号: **AYM30051**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	290kDa/292kDa
实测分子量	292kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,U-251MG,C6,Mouse brain
细胞定位	cilium,cytoplasm,cytoskeleton,cytosol,nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human USP9x.
序列	MTATTRGSPVGGNDNQGQAPDGQSQPPLQQNTSSPDSSNENSPATPPDEQGQGDAPPQLEDEEPAFPHTDLAKLDD MINRPRWVVPVLPKGEVLLEA

靶点信息

研究背景	This gene is a member of the peptidase C19 family and encodes a protein that is similar to ubiquitin-specific proteases. Though this gene is located on the X chromosome, it escapes X-inactivation. Mutations in this gene have been associated with Turner syndrome. Alternate transcriptional splice variants, encoding different isoforms, have been characterized.
------	---

基因ID	8239
基因名	USP9X
Swiss	Q93008
别名	USP9X;DFFRX;FAF;FAM;MRX99;MRXS99F;X-linked

产品验证



Western blot analysis of USP9x expressed in Mouse brain,293T,U-251MG using USP9x Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn